



UNIVERSIDAD NACIONAL  
AUTÓNOMA DE  
MÉXICO

## CAPÍTULO 7

### REFERENCIAS CITADAS

#### 7 Referencias

## 7. Referencias citadas

- Aguayo-Camargo, J.E., Marín-Córdova, S., Sánchez-Díaz, F., 1989. Evolución geológica de la Cuenca de México. Boletín de la Sociedad Mexicana de Mecánica de suelos. Tópicos geológicos de la Cuenca del Valle de México. Memoria de Simposio pags. 25-41.
- Al-Adamat, R., 2003. Groundwater vulnerability and risk mapping for the Basaltic aquifer of the Azraq basin of Jordan using GIS, Remote sensing and DRASTIC. *Applied Geography*, 23(4), pp.303-324.
- Aller, L., Bennett, T., Lehr, J.H., Petty, R.J., Hackett, G., 1987. DRASTIC: A standardized system for evaluating groundwater pollution potential using hydrogeologic settings. Environmental Protection Agency Report 600/2-87-035.
- Al-Zabet, T., 2002. Evaluation of aquifer vulnerability to contamination potential using the DRASTIC method. *Environmental Geology*, 43(1-2), pp.203- 208.
- Ahmed, A.A., 2009. Using Generic and Pesticide DRASTIC GIS-based models for vulnerability assessment of the Quaternary aquifer at Sohag, Egypt. *Hydrogeology Journal*, 17(5), 1203-1217.
- ASTER Digital Elevation Model, Land Processes Distributed Active Archive Center (LP DAAC), 2006.
- Auge M., 2004. Vulnerabilidad de acuíferos, conceptos y métodos. Universidad de Buenos Aires. CONICET. 2004.
- Babiker, I.S. et al., 2005. A GIS-based DRASTIC model for assessing aquifer vulnerability in Kakamigahara Heights, Gifu Prefecture, central Japan. *The Science of the total environment*, 345(1-3), pp.127-40.
- Cabral-Cano, E., H. Dixon, T., Miralles-Wilhelm, F., Díaz-Molina, O., Sánchez-Zamora, O. y E. Carande, R., 2008. Space geodetic imaging of rapid ground subsidence in Mexico City Geological Society of America, *Bull.*2008; 120: 1556-1566
- Denny, S.C., Allen, D.M. & Journeyay, J.M., 2007. DRASTIC-Fm: a modified vulnerability mapping method for structurally controlled aquifers in the southern Gulf Islands, British Columbia, Canada. *Hydrogeology Journal*, 15(3), pp.483-493.
- Ducci, D., 2000. GIS Techniques for Mapping Groundwater Contamination Risk. *Natural Hazards*, (1956), pp.279-294.

Durazo, J., 1996. Ciudad de México. Acuitardo superficial y contaminación acuífera. Ingeniería Hidráulica en México. Vol. XI, Núm 2, págs. 5-14.

ESRI ArcMap 9.3.1, ArcGIS Desktop 9.3.1, 2009.

Expósito, J.L. et al., 2010. Groundwater Protection Using Vulnerability Maps and Wellhead Protection Area (WHPA): A Case Study in Mexico. Water Resources Management, 24(15), pp.4219-4236.

FAO-UNESCO, 1974. Soil Map of the World (1:5,000,000), Vol. 1: Legend. UNESCO, Paris.

Ferrari, L., 2000. Avances en el conocimiento de la Faja Volcánica Transmexicana durante la última década. Boletín de la Sociedad Geológica Mexicana, V. LIII, 84-92.

Fritch, T.G. et al., 2000. An Aquifer Vulnerability Assessment of the Paluxy Aquifer, Central Texas, USA, Using GIS and a Modified DRASTIC Approach. Environmental Management, 25(3), pp.337-345.

Global Mapper Software LLC, 2010. Global Mapper 10.02

Gogu, R.C. & Dassargues, A., 2000. Current trends and future challenges in groundwater vulnerability assessment using overlay and index methods. Environmental Geology, 39(6), pp.549-559.

Gogu, R.C., Hallet, V. & Dassargues, A., 2003. Comparison of aquifer vulnerability assessment techniques. Application to the Néblon river basin (Belgium). Environmental Geology, 44(8), pp.881-892.

Golden Software, 2011. SURFER Version 10.1.561 (32-bit) Surface Mapping System.

Gómez-Tuena, A., Orozco-Esquivel, M. T., Ferrari, L., 2005. Petrogénesis ígnea de la Faja Volcánica Transmexicana. Boletín de la Sociedad Geológica Mexicana Volumen Conmemorativo del Centenario Temas Selectos de la Geología Mexicana Tomo LVII, núm. 3, 2005, p. 227-283

Jamrah, A., Al-Futaisi, A., Rajmohan, N., Al-Yaroubi, S., 2008. Assessment of groundwater vulnerability in the coastal region of Oman using DRASTIC index method in GIS environment. Environmental monitoring and assessment, 147(1-3), 125-38.

Javadi, S. et al., 2011. Modification of DRASTIC Model to Map Groundwater Vulnerability to Pollution Using Nitrate Measurements in Agricultural Areas., 13, pp.239-249.

Kim, Y.J., Hamm, S.Y. (1999). Assessment of the potential for groundwater contamination using the DRASTIC/EGIS technique, Cheongju area, South Korea. *Hydrogeology Journal*, 7, 227-235

Leyva, E., 2010. Acuíferos Semiconfinados y su modelacion: Aplicaciones al acuífero de la zona metropolitana de la Ciudad de México.

Liggett, J.E. & Allen, D.M., 2010. Evaluating the sensitivity of DRASTIC using different data sources, interpretations and mapping approaches. *Environmental Earth Sciences*, 62(8), pp.1577-1595.

MATLAB 7.10.0.499, R2010a. Interactive environment for algorithm development, data visualization, data analysis and numeric computation. The Mathworks Team, Natick, MA, USA. <http://mathworks.com>.

Mesri, G., Rokhsar, A., Bohor, B. F., 1975. Composition and compressibility of typical samples of Mexico City clay. *Géotechnique*, Volume 25, Issue 3, pages 527 –554, ISSN: 0016-8505, E-ISSN: 1751-7656

Mooser F., Molina T.C., 1992. Nuevo Modelo Hidrogeológico para la Cuenca de México. Centro de Investigación Sísmica Fundación Javier Barros Sierra, A.C. México. D.F., 19 pp.

Ormeci Varol, S. & Davraz, A., 2010. Hydrogeological investigation of Sarkikaraagac Basin (Isparta, Turkey) and groundwater vulnerability. *Water International*, 35(2), pp.177-194.

Ramos Leal, J. A. (2010a). Curso de vulnerabilidad acuífera e índices de calidad del agua. Geociencias.

Ramos Leal, J. A., Noyola Medrano, C., & Tapia Silva F. O. (2010b, February). Aquifer vulnerability and groundwater quality in mega cities: case of the Mexico Basin. *Environmental Earth Sciences*. *Environmental Earth Sciences*. 2010; 61(6):1309-1320.

Rushton, K.R., Rathod, K.S. Causes of non-linear step pumping test responses. *Q. J. Eng. Geol.* (1988) 21:147–158.

Santoyo Villa, E., Ovando-Shelly, E., Mooser H., F., 2005. Síntesis geotécnica de la Cuenca del Valle de México; TGC.

Shouyu, C. & Guangtao, F., 2003. A DRASTIC-based fuzzy pattern recognition methodology for groundwater vulnerability evaluation. *Hydrological Sciences Journal*, 48(2), pp.211-220.

Sistema de aguas de la Ciudad de México, 2005.

Sorichetta, A., 2011. Groundwater vulnerability assessment using statistical methods. Tesis de doctorado, UNIVERSITA DEGLI STUDI DI MILANO.

Tilahun, K. & Merkel, B.J., 2009. Assessment of groundwater vulnerability to pollution in Dire Dawa, Ethiopia using DRASTIC. Environmental Earth Sciences, 59(7), pp.1485-1496.

Vázquez Sánchez E., 1995. Modelo Conceptual hidrogeológico y características hidráulicas del acuífero en explotación en la parte meridional de la cuenca de México. México D.F.

Vazquez-Sánchez, E., Jaimes-Palomera, R., 1987. Geología de la Cuenca de México. Geofísica Internacional.

Veytia, B. M., 1975. Secretaría de Recursos Hídricos.

Voigt, H.-jürgen et al., 2004. Characterization of groundwater vulnerability to fulfill requirements of the water framework directive of the European Union. Water, 43, pp.567-574.